

Discrete Distributions

Now consider a series of *independent* experiments, each of which produces a Bernoulli random variable with probability of success p (p is the same for all of the trials)

Discrete Distributions

Now consider a series of *independent* experiments, each of which produces a Bernoulli random variable with probability of success p (p is the same for all of the trials)

The following discrete probability distributions arise from this model:

If the number of trials n is fixed in advance, the number of successes X has a **binomial** distribution

Discrete Distributions

Now consider a series of *independent* experiments, each of which produces a Bernoulli random variable with probability of success p (p is the same for all of the trials)

The following discrete probability distributions arise from this model:

If the number of trials n is fixed in advance, the number of successes X has a **binomial** distribution

If trials continue indefinitely until the first success is obtained, the number of failures obtained X has a **geometric** distribution.

Discrete Distributions

Now consider a series of *independent* experiments, each of which produces a Bernoulli random variable with probability of success p (p is the same for all of the trials)

The following discrete probability distributions arise from this model:

If the number of trials n is fixed in advance, the number of successes X has a **binomial** distribution

If trials continue indefinitely until the first success is obtained, the number of failures obtained X has a **geometric** distribution.

If trials continue indefinitely until the r^{th} success is obtained, the number of failures obtained X has a **negative binomial** distribution.

The Poisson Distribution

The probability mass function is:

$$f(x) = nb(x; r, p) = \binom{x + r - 1}{r - 1} p^r (1 - p)^x \quad x = 0, 1, 2, 3, \dots$$

The Poisson Distribution

The probability mass function is:

$$f(x) = nb(x; r, p) = \binom{x + r - 1}{r - 1} p^r (1 - p)^x \quad x = 0, 1, 2, 3, \dots$$

$$E(X) = \frac{r(1 - p)}{p} \quad V(X) = \frac{r(1 - p)}{p^2}$$

The Poisson Distribution

The probability mass function is:

$$f(x) = nb(x; r, p) = \binom{x+r-1}{r-1} p^r (1-p)^x \quad x = 0, 1, 2, 3, \dots$$

$$E(X) = \frac{r(1-p)}{p} \quad V(X) = \frac{r(1-p)}{p^2}$$

Computation:

Value	R	Spreadsheet
$P(X = x)$	$dnbinom(x, r, p)$	$= NEGBINOMDIST(x, r, p)$
$P(X \leq x)$	$pnbinom(x, r, p)$	

The Negative Binomial Distribution

Now we will perform some numerical experiments.

First generate a sample of 1,000,000 observations for a negative binomial experiment with $r = 3$ and probability of success $p = 0.4$:

```
x <- rnbinom(1000000, 3, 0.4)
```


The Negative Binomial Distribution

Now we will perform some numerical experiments.

First generate a sample of 1,000,000 observations for a negative binomial experiment with $r = 3$ and probability of success $p = 0.4$:

```
x <- rnbinom(1000000, 3, 0.4)
```

Now plot a histogram of the results:

```
hist(x)
```

The Negative Binomial Distribution

Now we will perform some numerical experiments.

First generate a sample of 1,000,000 observations for a negative binomial experiment with $r = 3$ and probability of success $p = 0.4$:

```
x <- rnbinom(1000000, 3, 0.4)
```

Now plot a histogram of the results:

```
hist(x)
```

To get a table of the results enter

```
table(x)
```

The Negative Binomial Distribution

Now we will perform some numerical experiments.

First generate a sample of 1,000,000 observations for a negative binomial experiment with $r = 3$ and probability of success $p = 0.4$:

```
x <- rnbinom(1000000, 3, 0.4)
```

Now plot a histogram of the results:

```
hist(x)
```

To get a table of the results enter

```
table(x)
```

The results through $X = 6$ should look something like:

0	1	2	3	4	5	6
63784	115545	138570	138259	124481	103801	8349

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Now compare the frequencies to the probabilities.

First compute the probability that $X = 0$:

dnbinom(0,3,0.4)

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Now compare the frequencies to the probabilities.

First compute the probability that $X = 0$:

```
dnbinom(0,3,0.4)
```

The result should be something like

```
[1] 0.064
```

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Now compare the frequencies to the probabilities.

First compute the probability that $X = 0$:

```
dnbinom(0,3,0.4)
```

The result should be something like

```
[1] 0.064
```

To get the probability that $X = 1$ enter

```
dnbinom(1,3,0.4)
```

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Now compare the frequencies to the probabilities.

First compute the probability that $X = 0$:

```
dnbinom(0,3,0.4)
```

The result should be something like

```
[1] 0.064
```

To get the probability that $X = 1$ enter

```
dnbinom(1,3,0.4)
```

This time the results should look something like:

```
[1] 0.1152
```

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Next compute the probability that $X = 2$:

dnbinom(2,3,0.4)

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Next compute the probability that $X = 2$:

```
dnbinom(2,3,0.4)
```

The result should be something like

```
[1] 0.13824
```

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Next compute the probability that $X = 2$:

```
dnbinom(2,3,0.4)
```

The result should be something like

```
[1] 0.13824
```

To get the probability that $X = 5$ enter

```
dnbinom(5,3,0.4)
```

The Negative Binomial Distribution

0	1	2	3	4	5	
63784	115545	138570	138259	124481	103801	8349

Next compute the probability that $X = 2$:

```
dnbinom(2,3,0.4)
```

The result should be something like

```
[1] 0.13824
```

To get the probability that $X = 5$ enter

```
dnbinom(5,3,0.4)
```

This time the results should look something like:

```
[1] 0.10451
```

The Negative Binomial Distribution

The expected value $E(X)$ in this case is:

$$E(X) = \frac{r(1-p)}{p} = \frac{3(.6)}{.4} = 4.5$$

The Negative Binomial Distribution

The expected value $E(X)$ in this case is:

$$E(X) = \frac{r(1-p)}{p} = \frac{3(.6)}{.4} = 4.5$$

To compute the sample mean \bar{x} , enter
mean(x)

The Negative Binomial Distribution

The expected value $E(X)$ in this case is:

$$E(X) = \frac{r(1-p)}{p} = \frac{3(.6)}{.4} = 4.5$$

To compute the sample mean \bar{x} , enter

mean(x) The result should be something like

[1] 4.499121

The Negative Binomial Distribution

The variance $V(X)$ in this case is:

$$V(X) = \frac{r(1-p)}{p^2} = \frac{3(.6)}{.4^2} = 11.25$$

The Negative Binomial Distribution

The variance $V(X)$ in this case is:

$$V(X) = \frac{r(1-p)}{p^2} = \frac{3(.6)}{.4^2} = 11.25$$

To compute the sample variance s^2 , enter
var(x)

The Negative Binomial Distribution

The variance $V(X)$ in this case is:

$$V(X) = \frac{r(1-p)}{p^2} = \frac{3(.6)}{.4^2} = 11.25$$

To compute the sample variance s^2 , enter `var(x)` The result should be something like

`[1] 11.2477`

The Negative Binomial Distribution

A fair coin is tossed until the second heads comes up.

Find the probability that the second heads comes up on the fifth toss ($x=3$).

The Negative Binomial Distribution

A fair coin is tossed until the second heads comes up.

Find the probability that the second heads comes up on the fifth toss ($x=3$).

Solution: 0.125

The Negative Binomial Distribution

A fair coin is tossed until the second heads comes up.

Find the probability that the second heads comes up on the fifth toss ($x=3$).

Solution: 0.125

$dnbinom(3, 2, 0.5)$

The Geometric Distribution

A fair coin is tossed until the fourth heads comes up.

Find the probability that the fourth heads comes up on the seventh toss or sooner $x \leq 3$.

The Geometric Distribution

A fair coin is tossed until the fourth heads comes up.

Find the probability that the fourth heads comes up on the seventh toss or sooner $x \leq 3$.

Solution: 0.5

The Geometric Distribution

A fair coin is tossed until the fourth heads comes up.

Find the probability that the fourth heads comes up on the seventh toss or sooner $x \leq 3$.

Solution: 0.5

pnbinom(3, 4, 0.5)

The Negative Binomial Distribution

A fair coin is tossed until the fifth heads comes up.

Find the probability that this takes more than 8 tosses
($x > 3$)

The Negative Binomial Distribution

A fair coin is tossed until the fifth heads comes up.

Find the probability that this takes more than 8 tosses
($x > 3$)

Solution: 0.63672

The Negative Binomial Distribution

A fair coin is tossed until the fifth heads comes up.

Find the probability that this takes more than 8 tosses
($x > 3$)

Solution: 0.63672

$$1 - \text{pnbinom}(3, 5, 0.5)$$

The Negative Binomial Distribution

A fair coin is tossed until the third heads comes up.

Find the probability that this takes 9 or more tosses ($x > 5$)

The Negative Binomial Distribution

A fair coin is tossed until the third heads comes up.

Find the probability that this takes 9 or more tosses ($x > 5$)

Solution: 0.22656

The Negative Binomial Distribution

A fair coin is tossed until the third heads comes up.

Find the probability that this takes 9 or more tosses ($x > 5$)

Solution: 0.22656

$$1 - \text{pnbinom}(4, 3, 0.5) \quad \text{or} \quad = 1 - \text{GEOMDIST}(7, 0.5, \text{TRUE})$$

(Spreadsheet function is for GNUMERIC. EXCEL does not have this function)

The Negative Binomial Distribution

A baseball player has a .300 batting average.

Find the probability that their second hit in a game occurs on the 5th time at bat. ($x = 3$)

The Negative Binomial Distribution

A baseball player has a .300 batting average.

Find the probability that their second hit in a game occurs on the 5th time at bat. ($x = 3$)

Solution: 0.12348

The Negative Binomial Distribution

A baseball player has a .300 batting average.

Find the probability that their second hit in a game occurs on the 5th time at bat. ($x = 3$)

Solution: 0.12348

dnbinom(3, 2, 0.3)