

The Binomial Distribution

Gene Quinn

Binomial Distribution

Recall that a **Bernoulli trial** is an experiment which has two outcomes, usually called *success* and *failure*.

Define a random variable X to be 1 if the outcome of the Bernoulli trial is success, and 0 if it is failure.

Binomial Distribution

Recall that a **Bernoulli trial** is an experiment which has two outcomes, usually called *success* and *failure*.

Define a random variable X to be 1 if the outcome of the Bernoulli trial is success, and 0 if it is failure.

Consider an experiment that consists of n independent Bernoulli trials, each with probability of success p .

Let X_i be the value of the random variable associated with the i^{th} Bernoulli trial, and define a new random variable Y by:

$$Y = \sum_{i=1}^n X_i$$

Binomial Distribution

Recall that a **Bernoulli trial** is an experiment which has two outcomes, usually called *success* and *failure*.

Define a random variable X to be 1 if the outcome of the Bernoulli trial is success, and 0 if it is failure.

Consider an experiment that consists of n independent Bernoulli trials, each with probability of success p .

Let X_i be the value of the random variable associated with the i^{th} Bernoulli trial, and define a new random variable Y by:

$$Y = \sum_{i=1}^n X_i$$

The random variable Y will assume values in the set $\{0, 1, \dots, n\}$.